

ABSTRACTS

XXth Belgian Week of Gastroenterology 2008

REDUCED COX-RELATED PRODUCTION OF VASOCONSTRICTORS AND NOT NO-MEDIATED VASODILATION IS A COMMON MECHANISM OF VASCULAR HYPOREACTIVITY IN DIFFERENT MODELS OF

VIRAL KINETICS CAN QUICKLY PREDICT VIROLOGICAL RESPONSE IN HCV PATIENTS WITH NORMAL ALT TREATED WITH PEGYLATED IFN ALPHA2B AND RIBAVIRIN : A PROSPECTIVE STUDY. P. Deltenre (1), V. Canva (2), M. El Nadi (2), A. Louvet (2), S. Dharancy (2), L. Bocket (2), A. Hollebecque (2), H. Castel (2),

CORRELATION BETWEEN HEPATIC PROGENITOR CELLS AND FIBROSIS IN THE CHOLINE DEFICIENT/ETHIONINE INDUCED MODEL OF CHRONIC LIVER INJURY. N. Van Hul, J. Abarca Quinones, Y. Horsmans, L. Isabelle. UCL Saint-Luc.

THERAPEUTIC EFFECTS OF VITAMINS A AND C ON EXPERIMENTAL LIVER DISEASE INDUCED BY A COMBINATION OF CCL4 AND PENTOBARBITAL IN RATS. S. Wamutu (1), S. Francque (1), S. Chatterjee (2),

Invited lecture

IMPACT OF AMP-ACTIVATED PROTEIN KINASE (AMPK) ON THE CONTROL OF HEPATIC STELLATE CELLS (HSCS) ACTIVATION IN VITRO. A. Da Silva (1), J. Abarca-Quinones (1), L. Hue (2), B. Violet (3), Y. Horsmans (1), I.A. Leclercq (1). (1) UCL, (2) ICP, (3) INSERM Paris, France.

Background and aims

Invited lecture

- D07 -

HEPATITIS B REACTIVATION IN IMMUNE SUPPRESSED PATIENTS AT RISK. D. Shouval. Jerusalem, Israël.

SURVIVAL OF PATIENTS CANDIDATES FOR INTESTINAL TRANSPLANTATION : A 3 YEARS PROSPECTIVE

TRANSIENT ELASTOGRAPHY VERSUS BIOCHEMICAL MARKERS FOR THE NONINVASIVE DIAGNOSIS

A META-ANALYSIS OF THREE GWA SCANS FOR CROHN'S DISEASE IDENTIFIES MULTIPLE NEW SUSCEPTIBILITY LOCI CONTRIBUTING TO A BETTER REPRESENTATION OF THE GENETIC ARCHITECTURE OF THE DISEASE. C. Libioulle. ULg.

Introduction and aim

EFFICACY OF INFLIXIMAB IN REFRACTORY POUCHITIS AND CD-RELATED COMPLICATIONS OF THE

POSTERS

- D49 -

SURGICAL TECHNIQUES AND COMPLICATIONS OF INTESTINE, COMBINED LIVER-INTESTINE AND FULL MULTIVISCERAL SPLANCHNIC TRANSPLANTATION (INCLUDING THE USE OF LIVE DONOR). J. Pirenne (1), D. Monbaliu (1), W. Coosemans (1), R. Aerts (1), G. Maleux (1), E. Benedetti (2). (1) KUL, (2) University Illinois Chicago, USA.

COVERED STENT TIPS IS ASSOCIATED WITH A BETTER SURVIVAL IN CIRRHOTIC PATIENTS WITH

Marker selection in colon tissue : Marker candidates identified by re-expression were screened using 37 real-time

Using the EORTC Quality of life Dysphagia score, revealed a clear, but non significant trend favouring SMA for solids being significant for semisolids ($p = 0.003$).

Conclusions

ABSTRACTS READ BY TITLE

- D77 -

THE DEGREE OF POLYMERISATION OF ARABINOXYLO-OLIGOSACCHARIDES PLAYS A ROLE TO

CASE-REPORT : CODFISH-ALLERGY-INDUCED OESOPHAGITIS. G. Vanneste (1), G. Lauwers (2),

MACROPHAGES AND HISTIOCYTES IN THE COLON AND DIARRHOEA ! WHICH DIAGNOSIS SHOULD YOU CONSIDER ? K. Deraedt, K. Geboes. KULeuven.

Introduction : Collections of foamy macrophages are a common finding in mucosal biopsies of the colon. They most-

P14ARF HYPERMETHYLATION PROFILE : NO CORRELATION WITH MICROSATELLITE INSTABILITY PHENOTYPE IN COLORECTAL CANCER. C. Nyiraneza, A. Kartheuser, R. Detry, C. Sempoux, K. Dahan. UCL Saint-Luc.

Introduction : *p14ARF*

COMPLETE METABOLIC RESPONSE WITH FDG DOES NOT INDICATE COMPLETE PATHOLOGIC

References

1. Leach *et al.*, *British J. Cancer*, 2005.
2. Tofts *et al.*, *J. Magn. Reson. Imaging*, 1999.
3. Larsson *et al.*, *Magn. Reson. Med.*, 1996.
4. St Lawrence *et al.*,

INTER-OBSERVER RELIABILITY OF DIFFERENT CT SCORING SYSTEMS IN ACUTE PANCREATITISSE(VELITYP

CROHN'S DISEASE PHENOTYPE, PROGNOSIS, AND LONG-TERM COMPLICATIONS : WHAT TO EXPECT ?
J. Cosnes. Pa

A PECULIAR CASE OF PANCREATITIS : BEWARE OF PERFORMING THE OBVIOUS ERCP. W. Laleman,

